arXiv:0705.2078v2 [math.GT] 28 May 2007

The abelianization of a symmetric mapping class group

Masatoshi Sato

Abstract

Let $\Sigma_{g,r}$ be a compact oriented surface of genus g with r boundary components. We determine the abelianization of the symmetric mapping class group $\hat{\mathcal{M}}_{(g,r)}(p_2)$ of a double unbranched cover $p_2 : \Sigma_{2g-1,2r} \to \Sigma_{g,r}$ using the Riemann constant, Schottky theta constant, and the theta multiplier. We also give lower bounds of the abelianizations of some finite index subgroups of the mapping class group.

Contents

0 Introduction **2**

1 The symmetric mapping class group **4**
1.1 Definition of the symmetric mapping class group 4
1.2 The action of the mapping class group on the equivalent classes of G-covers 5

2 A lower bound of the order of the cyclic group $H_1(\hat{\mathcal{M}}_{(g,r)}(p_2);\mathbf{Z})$ **7**
2.1 The first homology group $H_1(\Gamma_g(p_2);\mathbf{Z})$ 7
2.2 The coinvariant $H_1(\mathcal{I}_{g,r};\mathbf{Z})_{\mathcal{M}_{g,r}(p_2)}$ 11

3 A surjective homomorphism $\hat{\mathcal{M}}_{(g)}(p_2) \to \mathbf{Z}/4\mathbf{Z}$ **15**
3.1 The Jacobi variety and the Prym variety 15
3.2 Definition of the homomorphism $e : \hat{\mathcal{M}}_g(p_2) \to <\sqrt{-1}>$ 17
3.3 Proof of the main theorem 18

0 Introduction

Let g be a positive integer, $r \geq 0$, and S a set of n points in the interior of $\Sigma_{g,r}$. We denote by $\mathrm{Diff}_+(\Sigma_{g,r}, \partial\Sigma_{g,r}, S)$ the group of all orientation preserving diffeomorphisms which fix the boundary $\partial\Sigma_{g,r}$ pointwise, and map S onto itself allowing to permute it. The mapping class group $\mathcal{M}_{g,r}^n$ is the group of all isotopy classes $\pi_0 \mathrm{Diff}_+(\Sigma_{g,r}, \partial\Sigma_{g,r}, S)$ of such diffeomorphisms. We write simply $\mathcal{M}_{g,r} := \mathcal{M}_{g,r}^0$ and $\mathcal{M}_g^n := \mathcal{M}_{g,0}^n$. The mapping class group and its finite index subgroups play an important role in low-dimensional topology, in the theory of Teichmüller spaces and in algebraic geometry. For example, the level d mapping class group $\mathcal{M}_{g,r}[d]$ is defined to be the finite index subgroup of $\mathcal{M}_{g,r}$ which acts trivially on $H_1(\Sigma_{g,r}; \mathbf{Z}/d\mathbf{Z})$ for $d > 0$. It arises as the orbifold fundamental group of the moduli space of genus g curves with level d structure.

To compute the abelianizations, or equivalently, the first integral homology groups of finite index subgroups is one of the important problems in the mapping class groups. The Torelli group $\mathcal{I}_{g,r}$ is the subgroup which acts trivially on $H_1(\Sigma_{g,r}; \mathbf{Z})$. McCarthy[16] proved that the first rational homology group of a finite index subgroup that includes the Torelli group vanishes for $r = n = 0$, and more generally, Hain[6] proved it for any $r \geq 0$, $n \geq 0$.

Theorem 0.1 (McCarthy, Hain). *Let $\mathcal{M}$ be a finite index subgroup of $\mathcal{M}_{g,r}^n$ that includes the Torelli group where $g \geq 3$, $r \geq 0$. Then*

$$H_1(\mathcal{M}; \mathbf{Q}) = 0.$$

This theorem gives us little information about $H_1(\mathcal{M}; \mathbf{Z})$ as a finite group. In fact, Farb raised the problem to compute the abelianizations of the subgroup $\mathcal{M}_{g,r}[d]$ in [3] Problem 5.23 p.43. In this paper, we confine ourselves to the case $r = 0$ or 1 when it is not specified. For a finite regular cover p on $\Sigma_{g,r}$, possibly branched, Birman-Hilden[2] defined the symmetric mapping class group $\hat{\mathcal{M}}_{(g,r)}(p)$. That is closely related to a finite index subgroup of the mapping class group. As stated in subsection 1.1, the symmetric mapping class group is a finite group extension of a certain finite index subgroup of the mapping class group. In particular, we will have $H_1(\hat{\mathcal{M}}_{(g,r)}(p); \mathbf{Q}) = 0$ for all abel covers p. But in general, the first integral homology groups of symmetric mapping class groups and finite index subgroups of $\mathcal{M}_{g,r}^n$ are unknown.

One of the finite index subgroups, the spin mapping class group is defined by the subgroup of the mapping class group that preserves a spin structures on the surface. Lee-Miller-Weintraub[15] made the surjective homomophism from the spin mapping class group to $\mathbf{Z}/4\mathbf{Z}$ using the theta multiplier. Harer[8] proved that this homomorphism is in fact an isomorphism.

In this paper, we determine the symmetric mapping class group $\hat{\mathcal{M}}_{(g,r)}(p_2)$ of an unbranched double cover $p_2 : \Sigma_{2g-1,2r} \to \Sigma_{g,r}$ using the Riemann theta constant, Schottky theta constant, and the theta multiplier. We also compute a certain finite index subgroup $\mathcal{M}_{g,r}(p_2)$ of the mapping class group. That is included in the level 2 mapping class group $\mathcal{M}_{g,r}[2]$.

If we fix the symplectic basis of $H_1(\Sigma_{g,r}; \mathbf{Z})$, the action of mapping class group $\mathcal{M}_{g,r}^n$ on $H_1(\Sigma_{g,r}; \mathbf{Z})$ induces the surjective homomorphism

$$\iota : \mathcal{M}_{g,r} \to Sp(2g; \mathbf{Z}),$$

where $Sp(2g;\mathbf{Z})$ is the symplectic group of rank $2g$. Denote the image of $\mathcal{M}_{g,r}(p_2)$ under ι by $\Gamma_g(p_2)$. We also denote the image $\iota(\mathcal{M}_{g,r}[d])$ by $\Gamma_g[d]$, that is equal to the kernel $\mathrm{Ker}(Sp(2g;\mathbf{Z}) \to Sp(2g;\mathbf{Z}/d\mathbf{Z}))$ of mod d reduction. The main theorem is as follows.

Theorem 0.2. *For $r = 0, 1$, when genus $g \geq 4$,*

$$H_1(\hat{\mathcal{M}}_{(g,r)}(p_2);\mathbf{Z}) \cong H_1(\mathcal{M}_{g,1}(p_2);\mathbf{Z}) \cong \mathbf{Z}/4\mathbf{Z},$$

$$H_1(\mathcal{M}_g(p_2);\mathbf{Z}) \cong \begin{cases} \mathbf{Z}/4\mathbf{Z}, & \textit{if } g : \textit{odd}, \\ \mathbf{Z}/2\mathbf{Z}, & \textit{if } g : \textit{even}, \end{cases}$$

$$H_1(\Gamma_g(p_2);\mathbf{Z}) \cong \mathbf{Z}/2\mathbf{Z}.$$

After proving the theorem, we state that the first homology groups of the level d mapping class group $H_1(\mathcal{M}_{g,1}[d];\mathbf{Z})$ have many elements of order 4 for any even integer d (Proposition 3.8).

In section 1, we define the symmetric mapping class group, and describe the relation to a finite index subgroup of the mapping class group. In section 2, we prove that the integral homology groups of $\hat{\mathcal{M}}_{(g,r)}(p_2)$ and $\mathcal{M}_{g,r}(p_2)$ are cyclic groups of order at most 4. We also have $H_1(\Gamma_g(p_2);\mathbf{Z}) \cong \mathbf{Z}/2\mathbf{Z}$. In section 3, we construct an isomorphism $H_1(\hat{\mathcal{M}}_{(g,r)}(p_2);\mathbf{Z}) \cong \mathbf{Z}/4\mathbf{Z}$ using the Schottky theta constant and the theta multiplier to complete the proof of theorem 0.2.

1 The symmetric mapping class group

In this section, we define the symmetric mapping class group following Birman-Hilden[2], and prove some properties. In particular, we describe $\mathcal{M}_{g,r}(p) = \operatorname{Im} P$ by means of the action of the mapping class group on the equivalent classes of the covers in Subsection 1.2. We will see that the groups $\hat{\mathcal{M}}_{(g,r)}(p_2)$ and $\mathcal{M}_{g,r}(p_2)$ do not depend on the choice of the double cover p_2 up to isomorphism.

1.1 Definition of the symmetric mapping class group

Birman-Hilden[2] defined the symmetric mapping class group of a regular cover $p : \Sigma_{g',r'} \to \Sigma_{g,r}$, possibly branched as follows. Denote the deck transformation group of the cover by $\operatorname{Deck}(p)$.

Definition 1.1. *Let $C(p)$ be the centralizer of the deck transformation $\operatorname{Deck}(p)$ in the diffeomorphism group $\operatorname{Diff}_+(\Sigma_{g',r'})$. The symmetric mapping class group of the cover p is defined by*

$$\hat{\mathcal{M}}_{(g,r)}(p) = \pi_0(C(p) \cap \operatorname{Diff}_+(\Sigma_{g',r'}, \partial\Sigma_{g',r'})).$$

Let $S \subset \Sigma_{g,r}$ be the branch set of the cover p. For $\hat{f} \in C(p) \cap \operatorname{Diff}_+(\Sigma_{g',r'}, \partial\Sigma_{g',r'})$, there exists a unique diffeomorphism $f \in \operatorname{Diff}_+(\Sigma_{g,r}, \partial\Sigma_{g,r}, S)$ such that the diagram

$$\begin{tikzcd}
\Sigma_{g',r'} \arrow[r, "\hat{f}"] \arrow[d, "p"'] & \Sigma_{g',r'} \arrow[d, "p"'] \\
\Sigma_{g,r} \arrow[r, "f"] & \Sigma_{g,r}
\end{tikzcd}$$

commutes. Note that f maps the branch set S into itself. The diffeomorphism $f \in \operatorname{Diff}_+(\Sigma_{g,r}, \partial\Sigma_{g,r}, S)$ is called the projection of $\hat{f} \in C(p) \cap \operatorname{Diff}_+(\Sigma_{g',r'}, \partial\Sigma_{g',r'})$. For $[\hat{f}], [\hat{g}] \in \hat{\mathcal{M}}_{(g,r)}(p)$ such that $[\hat{f}] = [\hat{g}]$, an isotopy between $\hat{f}$ and $\hat{g}$ induces the isotopy on the base space $\Sigma_{g,r}$ between the projections f and g. Hence we can define the homomorphism

$$\begin{array}{cccc} P: & \hat{\mathcal{M}}_{(g,r)}(p) & \to & \mathcal{M}^n_{g,r}, \\ & [\hat{f}] & \mapsto & [f] \end{array}$$

where $n \geq 0$ is the order of S. We denote the image $\operatorname{Im} P \subset \mathcal{M}^n_{g,r}$ by $\mathcal{M}_{g,r}(p)$. The kernel of P is included in the group of isotopy classes of all the deck transformations in $\hat{\mathcal{M}}_{(g,r)}(p_2)$. Since any deck transformations without identity do not fix the boundary pointwise, we have $\operatorname{Ker} P = id$ when $r = 1$. When $r = 0$, $\operatorname{Ker} P$ consists of the isotopy classes of all the deck transformations. In particular, $\operatorname{Ker} P$ is a finite group. Apply the Lyndon-Hochschild-Serre spectral sequence to the group extension

$$1 \to \operatorname{Ker} P \to \hat{\mathcal{M}}_{(g,r)}(p) \to \mathcal{M}_{g,r}(p) \to 0,$$

then we have

$$H_*(\hat{\mathcal{M}}_{(g,r)}(p); \mathbf{Q}) \cong H_*(\mathcal{M}_{g,r}(p); \mathbf{Q}).$$

1.2 The action of the mapping class group on the equivalent classes of G-covers

For a finite group G and a finite set S, denote all the surjective homomorphisms $\pi_1(\Sigma_{g,r} - S, *) \to G$ by $\operatorname{Surj}(\pi_1(\Sigma_{g,r} - S, *), G)$. The group G acts on this set by inner automorphism. Denote the quotient set by

$$m(G, *) := \operatorname{Surj}(\pi_1(\Sigma_{g,r} - S, *), G)/\operatorname{Inn} G.$$

For paths $l, l' : [0,1] \to \Sigma_{g,r} - S$ such that $l(0) = l'(1)$, we define $l \cdot l'$ to be the path obtained by traversing first l' and then l. For a path $l : [0,1] \to \Sigma_{g,r} - S$, we define a isomorphism l_* by

$$\begin{array}{cccc} l_* : & \pi_1(\Sigma_{g,r} - S, l(0)) & \to & \pi_1(\Sigma_{g,r} - S, l(1)). \\ & \gamma & \mapsto & l \cdot \gamma \cdot l^{-1} \end{array}$$

If we pick a path l from $*$ to $*'$, we have the isomorphism $l_* : \pi_1(\Sigma_{g,r} - S, *) \cong \pi_1(\Sigma_{g,r} - S, *')$. Hence we also have the isomorphism

$$m(G, *) = m(G, *').$$

It is easy to see that this isomorphism does not depend on the choice of l, hence we denote $m(G) := m(G, *)$.

The mapping class group $\mathcal{M}_{g,r}$ acts on the set $m(G)$. In fact the diffeomorphism $f \in \operatorname{Diff}_+(\Sigma_{g,r}, \partial\Sigma_{g,r}, S)$ induces the map

$$\begin{array}{ccc} m(G) & \to & m(G) \\ {[c]} & \mapsto & [cf_*]. \end{array}$$

Proposition 1.2. *Let $c : \pi_1(\Sigma_{g,r} - S, *) \to G$ denote the monodromy homomorphism of a branched or unbranched G-cover $p : \Sigma_{g',r'} \to \Sigma_{g,r}$, where S is the branch set. The stabilizer of $[c] \in m(G)$ is equal to $\mathcal{M}_{g,r}(p)$.*

Proof. Suppose $[f] \in \mathcal{M}^n_{g,r}$ be in the stablizer of $[c]$. Since $[cf_*] = [c]$, there exists a path l from $*$ to $f(*)$ such that

$$c(l_*^{-1} f(\gamma)) = c(\gamma), \text{ for } \gamma \in \pi_1(\Sigma_{g,r} - S, *).$$

In particular, we have

$$\operatorname{Ker}(c) = l_*^{-1} f_*(\operatorname{Ker} c).$$

Hence the covers p and fp are equivalent. Choose a lift $\hat{l}$ of l, then there exists $\hat{f} \in \operatorname{Diff}(\Sigma_{g',r'})$ such that

$$p\hat{f} = fp : \Sigma_{g',r'} \to \Sigma_{g,r}, \text{ and } \hat{f}(\hat{l}(0)) = \hat{l}(1).$$

Then we have

$$\hat{f} c(\gamma) \hat{f}^{-1} = c(l_*^{-1} f(\gamma)) = c(\gamma) \in \operatorname{Diff}_+ \Sigma_{g',r'}.$$

Hence $\hat{f}$ is in the centralizer $C(p)$ of the deck transformation group $\operatorname{Deck}(p)$. When $r = 1$, $\operatorname{Deck}(p)$ acts on $\pi_0(\partial\Sigma_{g',r'})$ transitively. It is easy to see that for any $\hat{f} \in C(p)$, there exist $t \in \operatorname{Deck}(p)$ such that $\hat{f}t$ acts trivially on $\pi_0(\partial\Sigma_{g',r'})$. Therefore, there exists $t \in \operatorname{Deck}(p)$ such that $\hat{f}t \in C(p) \cap \operatorname{Diff}_+(\Sigma_{g',r'}, \partial\Sigma_{g',r'})$ and $f = P([\hat{f}t])$.

Conversely, suppose $f = P(\hat{f}) \in \mathcal{M}_{g,r}(p)$. Choose a path $\hat{l}$ such that $\hat{f}(\hat{l}(0)) = \hat{l}(1)$. Denote the projection $l = p\hat{l}$, then we have

$$c(l_*^{-1} f(\gamma)) = \hat{f} c(\gamma) \hat{f}^{-1} = c(\gamma) \in \mathrm{Diff}_+ \Sigma_{g',r'}.$$

Hence we have $[c] = [cf_*]$. $\square$

Hence, $\mathcal{M}_{g,r}(p)$ is a finite index subgroup of the mapping class group. In particular, if p is an abel cover, $\mathcal{M}_{g,r}(p)$ includes the Torelli group. By Theorem 0.1, we have $H_1(\mathcal{M}_{g,r}(p); \mathbf{Q}) = 0$. Consider the double covers on $\Sigma_{g,r}$. The number of the equivalent classes of double unbranched covers on $\Sigma_{g,r}$ are $2^{2g} - 1$. Since the action of mapping class group $\mathcal{M}_{g,r}$ on $m(\mathbf{Z}/2\mathbf{Z})$ is transitive, the subgroup $\mathcal{M}_{g,r}(p_2)$ does not depend on the choice of the double cover p_2 up to conjugate. It is easy to see that $\hat{\mathcal{M}}_{(g,r)}(p_2)$ is also unique up to isomorphism.

2 A lower bound of the order of the cyclic group $H_1(\hat{\mathcal{M}}_{(g,r)}(p_2);\mathbf{Z})$

In this section we prove that the integral homology groups of $\hat{\mathcal{M}}_{(g,r)}(p_2)$ and $\mathcal{M}_{g,r}(p_2)$ are cyclic groups of order at most 4. We compute $H_1(\Gamma_g(p_2);\mathbf{Z})$ in Subsection 2.1 and $H_1(\mathcal{I}_{g,r};\mathbf{Z})_{\mathcal{M}_{g,r}(p_2)}$ in Subsection 2.2 to obtain the lower bound.

In subsection 1.2, we proved that the symmetric mapping class group $\hat{\mathcal{M}}_{(g,r)}(p_2)$ and $\mathcal{M}_{g,r}(p_2)$ do not depend on the choice of the unbranched double cover p_2 up to isomorphism. Hence we fix the unbranched double cover p_2 whose monodromy $c \in \mathrm{Hom}(\pi_1(\Sigma_{g,r});\mathbf{Z}/2\mathbf{Z}) \cong H^1(\Sigma_{g,r};\mathbf{Z}/2\mathbf{Z})$ is equal to the Poincaré dual of B_g in Figure 1.

2.1 The first homology group $H_1(\Gamma_g(p_2);\mathbf{Z})$

In this subsection, using the generators of $\Gamma_g[2]$ in Igusa[9], we prove that $H_1(\Gamma_g(p_2);\mathbf{Z})$ is a cyclic group of order 2. We also prove that $H_1(\hat{\mathcal{M}}_{(g,r)}(p_2);\mathbf{Z})$ and $H_1(\mathcal{M}_{g,r}(p_2);\mathbf{Z})$ are cyclic of order at most 4 when genus $g \geq 4$, using the $\mathcal{M}_{g,r}$ module structure of the abelianization of the Torelli group determined by Johnson[12]. In particular, we obtain $H_1(\mathcal{M}_g(p_2);\mathbf{Z}) \cong \mathbf{Z}/2\mathbf{Z}$ if genus $g \geq 4$ is even. In the next section, we complete the proof of Theorem 0.2.

We consider $\Sigma_{g,1} = \Sigma_g - D^2 \subset \Sigma_g$. Pick simple closed curves $\{A_i, B_i\}_{i=1}^g \subset \Sigma_{g,r}$ as shown in Figure 1. They give a symplectic basis of $H := H_1(\Sigma_{g,r};\mathbf{Z})$ which we denote by the same symbol $\{A_i, B_i\}_{i=1}^g$. The action of the mapping class group on $H_1(\Sigma_{g,r};\mathbf{Z})$ induces

$$\iota : \mathcal{M}_{g,r} \to \mathrm{Sp}(2g,\mathbf{Z}).$$

We denote the Dehn twist along the simple closed curve A_g by $a \in \mathcal{M}_{g,r}$. Let S be a subsurface in $\Sigma_{g,r}$ as shown in Figure 2 and denote their mapping class groups which fix the boundary pointwise by $\mathcal{M}_S$. The inclusion $S \to \Sigma_g$ induces a homomorphism

$$i_S : \mathcal{M}_S \to \mathcal{M}_g.$$



Figure 1:



Figure 2:

As in Introduction, we denote by $\iota : \mathcal{M}_g \to \mathrm{Sp}(2g;\mathbf{Z})$ the homomorphism defined by the action of $\mathcal{M}_g$ on the homology group H, and denote the ring of integral n-square matrices by $M(n;\mathbf{Z})$ for a positive integer n. It is easy to see that the image of $i_S(\mathcal{M}_S)$ under ι is

$$\iota(i_S(\mathcal{M}_S)) = \left\{ \sigma = \begin{pmatrix} \alpha' & {}^tv_1 & \beta' & 0 \\ 0 & 1 & 0 & 0 \\ \gamma' & {}^tv_2 & \delta' & 0 \\ v_3 & k & v_4 & 1 \end{pmatrix} \in \mathrm{Sp}(2g;\mathbf{Z}) \;\middle|\; \begin{matrix} \alpha',\beta',\gamma',\delta' \in M(g-1;\mathbf{Z}), \\ v_1,v_2,v_3,v_4 \in \mathbf{Z}^{g-1}, k \in \mathbf{Z} \end{matrix} \right\}.$$

Proposition 2.1. *When $g \geq 1$, $\Gamma_g(p_2)$ is generated by $\iota(i_S(\mathcal{M}_S))$ and $\iota(a^2)$.*

Proof. First, we show that $\Gamma_g(p_2)$ is generated by $\iota(i_S(\mathcal{M}_S))$ and $\Gamma_g[2]$. Since an element $\sigma \in \Gamma_g(p_2)$ preserves the homology class $B_g \in H_1(\Sigma_{g,r};\mathbf{Z}/2\mathbf{Z})$, it can be written in the form

$$\sigma \equiv \begin{pmatrix} \alpha' & {}^tv_1 & \beta' & 0 \\ 0 & 1 & 0 & 0 \\ \gamma' & {}^tv_2 & \delta' & 0 \\ v_3 & k & v_4 & 1 \end{pmatrix} \bmod 2.$$

Hence there exists $\sigma_0 \in \iota(i_S(\mathcal{M}_S))$ such that

$$\sigma_0 \equiv \sigma \mod 2,$$

so that $\Gamma_g(p_2)$ is generated by $\iota(i_S(\mathcal{M}_S))$ and $\Gamma_g[2]$.

Next, we describe the generators of $\Gamma_g[2]$ given in Igusa[9]. We denote by I_n the unit matrix of order n, and by e_{ij} the $2g$-square matrix with 1 at the (i,j)-th entry and 0 elsewhere. As was shown in Igusa[9], $\Gamma_g[2]$

is generated by

$$\begin{aligned}
\alpha_{ij} &= I_{2g} + 2e_{ij} - 2e_{g+j,g+i} && 1 \leq i, j \leq g,\ i \neq j, \\
\alpha_{ii} &= I_{2g} - 2e_{ii} - 2e_{i+g,i+g} && 1 \leq i \leq g, \\
\beta_{ij} &= I_{2g} + 2e_{i,j+g} + 2e_{j,i+g} && 1 \leq i < j \leq g, \\
\beta_{ii} &= I_{2g} + 2e_{i,i+g} && 1 \leq i \leq g, \\
\gamma_{ij} &= {}^{t}\beta_{ij} && 1 \leq i \leq j \leq g.
\end{aligned}$$

To prove the proposition, it suffices to show that these matrices are in the subgroup of $\Gamma_g(p_2)$ generated by $\iota(i_S(\mathcal{M}_S))$ and $\iota(a^2)$. The matrices

$$\alpha_{ij}(1 \leq i \leq g-1, 1 \leq j \leq g),\ \beta_{ij}(1 \leq i \leq j \leq g-1), \text{ and } \gamma_{ij}(1 \leq i \leq j \leq g)$$

are clearly in $\iota(i_S(\mathcal{M}_S))$. Choose oriented simple closed curves $C_i, C'_i, C_{ij}, C'_{ij}, C''_{ij} \subset \Sigma_{g,r}$ such that $[C_i] = A_i$, $[C'_i] = B_i$, $[C_{ij}] = A_i + A_j$, $[C'_{ij}] = B_i + B_j$, $[C''_{ij}] = A_i + B_j$. Denote the Dehn twist along a simple closed curve C by T_C. Then the matrices

$$\alpha_{gj}(1 \leq j \leq g-1) \ \text{ and } \ \beta_{ig}(1 \leq i \leq g-1)$$

are written as $\iota(T^2_{C''_{gi}} T^{-2}_{C'_i} T^{-2}_{C_g})$ and $\iota(T^2_{C_{ig}} T^{-2}_{C_i} T^{-2}_{C_g})$ respectively. Clearly $\iota(T^2_{C'_i})$ and $\iota(T^2_{C_i})$ are in $\iota(i_S(\mathcal{M}_S))$, and we have $\iota(T^2_{C_g}) = \iota(a^2)$. Denote the two boundary components of S by S_1 and S_2. For any two arcs $l_1, l_2 : [0,1] \to S$ that satisfy $l_1(0) = l_2(0) \in S_1$ and $l_1(1) = l_2(1) \in S_2$, there exists $\varphi \in \mathcal{M}_S$ such that

$$\varphi l_1 = l_2.$$

Choose C''_{gi} and C_{ig} such that $\sharp(C''_{gi} \cap S_1) = \sharp(C_{ig} \cap S_1) = 1$ and they intersect with S_1 transversely, there exist $\psi, \psi' \in i_S(\mathcal{M}_S)$ that satisfy $[\psi(C''_{gi})] = [\psi'(C_{ig})] = A_g$. Thus we have

$$\psi T^2_{C''_{gi}} \psi^{-1} = \psi' T^2_{C_{ig}} \psi'^{-1} = a^2.$$

This proves the matrices α_{gj} and β_{ig} are in the subgroup of $\Gamma_g(p_2)$ generated by $\iota(i_S(\mathcal{M}_S))$ and $\iota(a^2)$. Finally the matrices α_{gg} and β_{gg} satisfy $\alpha_{gg} = \iota(T^2_{C''_{gg}})\beta_{gg}\gamma_{gg}^{-1}$, and $\beta_{gg} = \iota(a^2)$. Hence α_{gg} and β_{gg} are also in the subgroup, as was to be shown. □

Using Proposition 2.1, we now calculate the first homology group $H_1(\Gamma_g(p_2); \mathbf{Z})$.

Proposition 2.2. *When* $g \geq 4$,

$$H_1(\Gamma_g(p_2); \mathbf{Z}) \cong \mathbf{Z}/2\mathbf{Z}.$$

Proof. Powell [17] had proved $H_1(\mathcal{M}_g; \mathbf{Z}) = 0$ when $g \geq 3$. More generally, Harer [7] proved that $H_1(\mathcal{M}_{g,r}; \mathbf{Z}) = 0$ when $g \geq 3$ for any r. Hence the first homology $H_1(\mathcal{M}_S; \mathbf{Z})$ vanishes since genus of $S \geq 3$. We have

$$i_S(\mathcal{M}_S) = \{0\} \subset H_1(\mathcal{M}_g(p_2); \mathbf{Z}).$$

Since we proved that the group $\Gamma_g(p_2)$ is generated by $\iota(i_S(\mathcal{M}_S))$ and $\iota(a^2)$ in Proposition 2.1, the homology group $H_1(\Gamma_g(p_2);\mathbf{Z})$ is generated by $[\iota(a^2)]$.

Next, we construct a surjective homomorphism $\Gamma_g(p_2) \to \mathbf{Z}/2\mathbf{Z}$. Since any $\sigma = (\sigma_{ij}) \in \Gamma_g(p_2)$ preserves the homology class $B_g \in H_1(\Sigma_{g,r};\mathbf{Z}/2\mathbf{Z})$, we have

$$\sigma_{gi} \equiv \delta_{ig}, \text{ and } \ \sigma_{i\,2g} \equiv \delta_{i\,2g} \mod 2,$$

where δ is the Kronecker delta. Then for $\sigma, \sigma' \in \Gamma_g(p_2)$, the $(g, 2g)$-th entry of $\sigma\sigma'$ satisfies

$$(\sigma\sigma')_{g\,2g} = \sum_{i=1}^{2g} \sigma_{gi}\sigma'_{i\,2g} \equiv \sigma_{g\,2g} + \sigma'_{g\,2g} \mod 4.$$

Hence we have the homomorphism

$$\begin{array}{cccc} \Psi: & \Gamma_g(p_2) & \to & \mathbf{Z}/2\mathbf{Z} \\ & \sigma & \mapsto & \dfrac{\sigma_{g\,2g}}{2}. \end{array}$$

Since $\Psi([\iota(a^2)]) = 1$, we have $[\iota(a^2)] \neq 0 \in H_1(\Gamma_g(p_2);\mathbf{Z})$.

Finally, to complete the proof it suffices to show that $2[\iota(a^2)] = 0$. Apply the Lyndon-Hochschild-Serre spectral sequence to the group extension

$$1 \to \mathcal{I}_{g,r} \to \mathcal{M}_{g,r}(p_2) \to \Gamma_g(p_2) \to 0,$$

then we have

$$H_1(\mathcal{I}_{g,r};\mathbf{Z})_{\mathcal{M}_{g,r}(p_2)} \to H_1(\mathcal{M}_{g,r}(p_2);\mathbf{Z}) \to H_1(\Gamma_g(p_2);\mathbf{Z}) \to 0.$$

Denote by D and D' the simple closed curves as shown in Figure 3. Denote by c_1, c_2, and c_3 the Dehn twists along the simple closed curves C_1, C_2, and C_3 as shown in Figure 4 respectively. Since c_1 and c_2 are in $i_S(\mathcal{M}_S)$, $[c_1] = [c_2] = 0 \in H_1(\mathcal{M}_{g,r}(p_2);\mathbf{Z})$. By the chain relation, we have $T_DT_{D'} = (c_1c_2c_3)^4$. Using the braid relations $c_1c_3 = c_3c_1$, $c_1c_2c_1 = c_2c_1c_2$, and $c_2c_3c_2 = c_3c_2c_3$, we have



Figure 3:

$$[T_DT_{D'}] = [(c_1c_2c_3)^4] = [c_3c_2c_1^2c_2c_3] = [c_3c_2c_1^2c_2^{-1}c_3^{-1}] + [c_3c_2^2c_3^{-1}] + [c_3^2] \in H_1(\mathcal{M}_{g,r}(p_2);\mathbf{Z}).$$



Figure 4:

Since $c_3c_2c_1^2c_2^{-1}c_3^{-1}$ and $c_2c_1^2c_2^{-1}$ are the squares of the Dehn twists along the simple closed curves $c_3c_2(C_1)$ and $c_3(C_2)$, we have

$$[c_3c_2c_1^2c_2^{-1}c_3^{-1}] = [c_3c_2^2c_3^{-1}] = [c_3^2] = [a^2].$$

Hence $[T_DT_{D'}^{-1}] = [T_DT_{D'}] + [T_{D'}^{-2}] = 2[a^2]$. Since $T_DT_{D'}^{-1} \in \mathcal{I}_{g,r}$, it follows that $2[\iota(a^2)] = [\iota(T_DT_{D'}^{-1})] = 0 \in H_1(\Gamma_g(p_2);\mathbf{Z})$. This proves the proposition. ☐

2.2 The coinvariant $H_1(\mathcal{I}_{g,r};\mathbf{Z})_{\mathcal{M}_{g,r}(p_2)}$

To calculate the first homology group of the symmetric mapping class groups, we compute $H_1(\mathcal{I}_{g,r};\mathbf{Z})_{\mathcal{M}_{g,r}(p_2)}$.

Lemma 2.3. *When $g \geq 4$,*

$$H_1(\mathcal{I}_g;\mathbf{Z})_{\mathcal{M}_g(p_2)} \cong \begin{cases} \mathbf{Z}/2\mathbf{Z}, & \text{if } g: \text{odd}, \\ 0, & \text{if } g: \text{even}, \end{cases}$$

$$H_1(\mathcal{I}_{g,1};\mathbf{Z})_{\mathcal{M}_{g,1}(p_2)} \cong \mathbf{Z}/2\mathbf{Z}.$$

Moreover $H_1(\mathcal{I}_{g,r};\mathbf{Z})_{\mathcal{M}_{g,r}(p_2)}$ is generated by $T_DT_{D'}^{-1} \in \mathcal{I}_{g,r}$ for $r = 0, 1$.

Before proving the lemma, we review the space of boolean polynomials. Let H denote the first homology group $H_1(\Sigma_{g,r};\mathbf{Z})$ of the surface as before. Consider the polynomial ring with coefficients in $\mathbf{Z}/2\mathbf{Z}$ with the basis $\bar{x}$ for $x \in H \otimes \mathbf{Z}/2\mathbf{Z}$. Denote by J the ideal in the polynomial generated by

$$\overline{x+y} - (\bar{x} + \bar{y} + x \cdot y), \quad \bar{x}^2 - \bar{x}, \qquad \text{for } x, y \in H \otimes \mathbf{Z}/2\mathbf{Z}.$$

The space of boolean polynomials of degree at most n is defined by

$$B^n = \frac{M_n}{J \cap M_n},$$

where M_n is the module of all polynomials of degree at most n. Note that B^n is isomorphic to the $\mathbf{Z}/2\mathbf{Z}$ module of all square free polynimials of degree at most n generated by $\{\bar{A}_i, \bar{B}_i\}_{i=1}^{g}$.

Denote B^3 by $B^3_{g,1}$, and for $\alpha = \Sigma_{i=1}^{g} \bar{A}_i \bar{B}_i \in B^2$, the cokernel of

$$\begin{array}{ccc} B^1 & \rightarrow & B^3 \\ x & \mapsto & \alpha x \end{array}$$

by $B^3_{g,0}$. The action of $\mathcal{M}_{g,r}$ on H induces an action on $B^3_{g,r}$. Birman-Craggs[1] defined a family of homomorphisms $\mathcal{I}_g \to \mathbf{Z}/2\mathbf{Z}$. Johnson[13] showed that these homomorphisms give a surjective homomorphism of $\mathcal{M}_{g,r}$ modules

$$\mu : \mathcal{I}_{g,r} \to B^3_{g,r}.$$

For $r = 0, 1$, Johnson[14] showed that the induced homomorphism $H_1(\mathcal{I}_{g,r}; \mathbf{Z})_{\mathcal{M}_{g,r}[2]} \cong B^3_{g,r}$ is an isomorphism.

proof of Lemma 2.3. Since μ is an isomorphism of $\mathcal{M}_{g,r}$ module, we have

$$H_1(\mathcal{I}_{g,r}; \mathbf{Z})_{\mathcal{M}_{g,r}(p_2)} \cong (B^3_{g,r})_{\mathcal{M}_{g,r}(p_2)}.$$

Hence it suffices to compute $(B^3_{g,r})_{\mathcal{M}_{g,r}(p_2)}$ to prove the lemma. Denote the subsurface $S' \subset S$ of genus $g-1$ as shown in Figure 2. $\mathcal{I}_{S'}$ is the Torelli group of S', that is the subgroup of $\mathcal{M}_{S'}$ which act trivially on $H_1(S'; \mathbf{Z})$. Consider the homomorphism

$$(\mathcal{I}_{S'})_{\mathcal{M}_{S'}} \to (\mathcal{I}_{g,r})_{\mathcal{M}_{g,r}(p_2)} \cong (B^3_{g,r})_{\mathcal{M}_{g,r}(p_2)}.$$

induced by the inclusion $S' \to \Sigma_{g,r}$. Since $(\mathcal{I}_{S'})_{\mathcal{M}_{S'}} = 0$ (Johnson[11]), the image of the homomorphism is trivial. Thus we have

$$\bar{1} = \bar{X} = \bar{X}\bar{Y} = \bar{X}\bar{Y}\bar{Z} = 0, \quad \text{for } \{X, Y, Z\} \subset \{A_1, A_2, \cdots, A_{g-1}, B_1, B_2 \cdots, B_{g-1}\}.$$

For $X = A_g, B_g$, we have

$$(I_{2g} + e_{1,g+1})(\bar{B}_1\bar{X}) = (\bar{B}_1 + \bar{A}_1 + 1)\bar{X}, \ (I_{2g} + e_{g+1,1})(\bar{A}_1\bar{X}) = (\bar{A}_1 + \bar{B}_1 + 1)\bar{X},$$
$$\text{and} \ \ (I_{2g} + e_{1,2} - e_{g+2,g+1})(\bar{A}_2\bar{X}) = (\bar{A}_2 + \bar{A}_1)\bar{X}.$$

Hence $\bar{X} = \bar{A}_1\bar{X} = \bar{B}_1\bar{X} = 0 \in (B^3_{g,r})_{\mathcal{M}_{g,r}(p_2)}$. For $1 < i < g$, we have

$$(I_{2g} + e_{g+i,1} + e_{g+1,i})(\bar{A}_1\bar{X}) = (\bar{A}_1 + \bar{B}_i)\bar{X}, \ \text{and} \ \ (I_{2g} + e_{i,g+1} + e_{1,g+i})(\bar{B}_1\bar{X}) = (\bar{B}_1 + \bar{A}_i)\bar{X}.$$

Hence $\bar{B}_i\bar{X} = \bar{A}_i\bar{X} = 0 \in (B^3_{g,r})_{\mathcal{M}_{g,r}(p_2)}$. If we put $\bar{X} = \bar{A}_g\bar{B}_g$, we have $\bar{Y}\bar{A}_g\bar{B}_g = 0$ in the same way for $\bar{Y} \in \{1, \bar{A}_1, \bar{A}_2, \cdots, \bar{A}_{g-1}, \bar{B}_1, \bar{B}_2, \cdots, \bar{B}_{g-1}\}$. For $X = A_g, B_g$, and any i, j such that $1 \leq i, j < g$, $i \neq j$, we have

$$(I_{2g} + e_{g+j,j})(\bar{A}_i\bar{A}_j\bar{X}) = \bar{A}_i\bar{A}_j\bar{X} + \bar{A}_i\bar{B}_j\bar{X} + \bar{A}_i\bar{X}, \ (I_{2g} + e_{j,g+j})(\bar{A}_i\bar{B}_j\bar{X}) = \bar{A}_i\bar{B}_j\bar{X} + \bar{A}_i\bar{A}_j\bar{X} + \bar{A}_i\bar{X},$$
$$(I_{2g} + e_{g+j,j})(\bar{B}_i\bar{A}_j\bar{X}) = \bar{B}_i\bar{A}_j\bar{X} + \bar{B}_i\bar{B}_j\bar{X} + \bar{B}_i\bar{X}, \ (I_{2g} + e_{g+i,g} + e_{2g,g+i})(\bar{A}_i\bar{A}_g\bar{B}_g) = \bar{A}_i\bar{A}_g\bar{B}_g + \bar{A}_i\bar{B}_i\bar{B}_g,$$
$$\text{and} \ \ (I_{2g} - e_{1,1} - e_{g+1,g+1} + e_{i,1} + e_{1,i} + e_{g+i,g+1} + e_{g+1,g+i})(\bar{A}_1\bar{B}_1\bar{A}_g) = \bar{A}_i\bar{B}_i\bar{A}_g.$$

Hence $\bar{A}_i\bar{B}_j\bar{X} = \bar{A}_i\bar{A}_j\bar{X} = \bar{B}_i\bar{B}_j\bar{X} = \bar{A}_i\bar{B}_i\bar{B}_g = 0$, and $\bar{A}_1\bar{B}_1\bar{A}_g = \bar{A}_i\bar{B}_i\bar{A}_g \in (B^3_{g,r})_{\mathcal{M}_{g,r}(p_2)}$.

Therefore $(B^3_{g,r})_{\mathcal{M}_{g,r}(p_2)}$ is a cyclic group of order 2 with generator $\bar{A}_1\bar{B}_1\bar{A}_g$ or a trivial group. For $r = 0$, $B^3_{g,0}$ has a relation

$$\alpha\bar{A}_g = (\sum_{i=1}^{g} \bar{A}_i\bar{B}_i)\bar{A}_g = 0,$$

so that $(g-1)\bar{A}_1\bar{B}_1\bar{A}_g = 0 \in (B^3_{g,0})_{\mathcal{M}_g(p_2)}$. This shows that $(B^3_{g,r})_{\mathcal{M}_{g,r}(p_2)}$ is trivial when g is even and $r = 0$.

Next we consider the case g is odd or $r = 1$. Let S_n be the permutation group of degree n and sign(s) the sign of $s \in S_n$. Denote by $\Lambda^n H$ the image of the homomorphism

$$\begin{array}{rccc} \lambda: & H^{\otimes n} & \to & H^{\otimes n} \\ & x_1 \otimes x_2 \otimes \cdots \otimes x_n & \mapsto & \displaystyle\sum_{s \in S_n} \mathrm{sign}(s) x_{s(1)} \otimes x_{s(2)} \otimes \cdots \otimes x_{s(n)}. \end{array}$$

Denote by V_1 and V_0 the module $\Lambda^3 H$ and the cokernel of

$$\begin{array}{ccc} H & \to & \Lambda^3 H \\ X & \mapsto & \sum_{i=1}^{g} A_i \wedge B_i \wedge X, \end{array}$$

respectively. Then Johnson[12] shows

$$\begin{array}{ccc} \dfrac{B^3_{g,r}}{B^2} & \to & V_r \otimes \mathbf{Z}/2\mathbf{Z} \\ \bar{X}\bar{Y}\bar{Z} & \mapsto & X \wedge Y \wedge Z, \end{array}$$

is a well-defined $\mathcal{M}_{g,r}$ module isomorphism. Now we have a $\mathcal{M}_{g,r}(p_2)$ homomorphism

$$\begin{array}{rccc} (B_g\cdot)C: & (B^3_{g,r})_{\mathcal{M}_{g,r}(p_2)} & \to & \mathbf{Z}/2\mathbf{Z} \\ & \bar{X}\bar{Y}\bar{Z} & \mapsto & (X\cdot Y)B_g\cdot Z + (Y\cdot Z)B_g\cdot X + (Z\cdot X)B_g\cdot Y. \end{array}$$

Here it should be remarked that the intersection number with B_g $(B_g\cdot) : H \otimes \mathbf{Z}/2\mathbf{Z} \to \mathbf{Z}/2\mathbf{Z}$ is $\mathcal{M}_{g,r}(p_2)$-invariant. Since $(B_g\cdot)C(\bar{A}_1\bar{B}_1\bar{A}_g) = 1$, it is surjective. Hence $(B^3_{g,r})_{\mathcal{M}_{g,r}(p_2)}$ is a cyclic group of order 2 with generator $\bar{A}_1\bar{B}_1\bar{A}_g$. Johnson[13] computed $\mu(T_D T_{D'}^{-1}) = \bar{A}_1\bar{B}_1(\bar{A}_g + 1)$, so that $T_D T_{D'}^{-1}$ is a generator of $H_1(\mathcal{I}_{g,r};\mathbf{Z})_{\mathcal{M}_{g,r}(p_2)}$. □

Now, we prove that $H_1(\hat{\mathcal{M}}_{(g,r)};\mathbf{Z})$ and $H_1(\mathcal{M}_{g,r};\mathbf{Z})$ are cyclic groups of order at most 4. We need the following Lemma.

Lemma 2.4. *Let $\hat{b} : \hat{\mathcal{M}}_{(g,1)}(p_2) \to \hat{\mathcal{M}}_{(g)}(p_2)$ be a homomorphism induced by an obvious embedding $\Sigma_{2g-1,2} \to \Sigma_{2g-1}$. Then $\hat{b}$ is surjective.*

Proof. By the obvious embedding $\Sigma_{g,1} \to \Sigma_g$, we have a surjective homomorphism $b : \mathcal{M}_{g,1}(p_2) \to \mathcal{M}_g(p_2)$. Since the diagram

$$\begin{tikzcd} \hat{\mathcal{M}}_{(g,1)}(p_2) \arrow[r, "\hat{b}"] \arrow[d, "P"'] & \hat{\mathcal{M}}_{(g)}(p_2) \arrow[d, "P"'] \\ \mathcal{M}_{g,1}(p_2) \arrow[r, "b"] & \mathcal{M}_g(p_2) \end{tikzcd}$$

commutes, $\hat{b}P = Pb$ is surjective. Hence it surfices to show $\operatorname{Ker} P \subset \hat{\mathcal{M}}_{(g)}(p_2)$ is included in $\operatorname{Im} \hat{b}$. Recall that $\operatorname{Ker} P$ consists of the isotopy classes of all the deck transformation. Cut the surface $\Sigma_{g,r}$ along the two simple closed curves A_g, A'_g in Figure 5. Then we have the subsurface S_0 of genus $g-1$ and the other subsurface S'_0 of genus 0. We can construct a diffeomorphism $\hat{f}_0 \in C(p) \cap \mathrm{Diff}_+(\Sigma_{2g-1,2}, \partial\Sigma_{2g-1,2})$ which have the following properties:



Figure 5:

(i) $\hat{f}_0|_{p^{-1}(S_0)}$ is the restriction of the deck transformation $t \neq id$,

(ii) $\hat{f}_0|_{p^{-1}(S_1)} = {T'}_{A_g} {T'}^{-1}_{A'_g}$, where ${T'}_{A_g}$ and ${T'}_{A'_g}$ is the half Dehn twists along A_g and A'_g.

Then $\hat{f}_0$ is included in $C(p) \cap \mathrm{Diff}_+(\Sigma_{2g-1,2}, \partial\Sigma_{2g-1,2})$, and the image of $[\hat{f}_0]$ under $\hat{\mathcal{M}}_{(g,1)}(p_2) \to \hat{\mathcal{M}}_{(g)}(p_2)$ equals the deck transformation t. This proves the lemma. ☐

In the proof of Proposition 2.2, we have the exact sequence

$$H_1(\mathcal{I}_{g,r}; \mathbf{Z})_{\mathcal{M}_{g,r}(p_2)} \to H_1(\mathcal{M}_{g,r}(p_2); \mathbf{Z}) \to H_1(\Gamma_g(p_2); \mathbf{Z}) \to 0.$$

By Proposition 2.2 and Lemma 2.3, we obtain

$$H_1(\mathcal{M}_{g,r}(p_2); \mathbf{Z}) = \mathbf{Z}/2\mathbf{Z} \text{ or } \mathbf{Z}/4\mathbf{Z}.$$

In particular if genus g is even,

$$H_1(\mathcal{M}_g(p_2); \mathbf{Z}) = \mathbf{Z}/2\mathbf{Z}.$$

From the isomorphism $\hat{\mathcal{M}}_{(g,1)}(p_2) \cong \mathcal{M}_{g,1}(p_2)$ and the surjective homomorphism $b : \hat{\mathcal{M}}_{(g,1)}(p_2) \to \hat{\mathcal{M}}_{(g)}(p_2)$, we have

$$H_1(\hat{\mathcal{M}}_{(g,r)}(p_2); \mathbf{Z}) = \mathbf{Z}/2\mathbf{Z} \text{ or } \mathbf{Z}/4\mathbf{Z}$$

for $r = 0, 1$.

Remark 2.5. *For $r = 0, 1$, pick a simple closed curve $c \subset \Sigma_{g,r}$. If the intersection number $c \cdot B_g$ is odd, then $[T_c^2] \in H_1(\Gamma_g(p_2); \mathbf{Z})$ is a generator. Hence $[T_c^2] \in H_1(\mathcal{M}_{g,r}(p_2); \mathbf{Z})$ is also a generator, and the lift of T_c^2 is a generator of $H_1(\hat{\mathcal{M}}_{(g,r)}(p_2); \mathbf{Z})$.*

If c is included in the subsurface S, we have $[T_c^2] = 0 \in H_1(\hat{\mathcal{M}}_{(g,r)}(p_2); \mathbf{Z})$, by Proposition 2.1.

3 A surjective homomorphism $\hat{\mathcal{M}}_{(g)}(p_2) \to \mathbf{Z}/4\mathbf{Z}$

For a root of unity ζ, we denote by $<\zeta>$ the cyclic group generated by ζ. In this section, we construct a surjective homomorphism

$$e : \hat{\mathcal{M}}_{(g)}(p_2) \to <\sqrt{-1}>$$

using the Schottky theta constant associated with the cover $p_2 : \Sigma_{2g-1} \to \Sigma_g$ when $g \geq 2$, to complete Theorem 0.2. In the following, suppose genus $g \geq 2$.

3.1 The Jacobi variety and the Prym variety

Endow the surface Σ_g with the structure of a Riemann surface R. Then the covering map $p_2 : \Sigma_{2g-1} \to \Sigma_g$ induces the structure of a Riemann surface $\hat{R}$ in the surface Σ_{2g-1}. In this subsection, we review the Jacobi variety of the Riemann surface R and the Prym variety of the double unbranched cover $p_2 : \hat{R} \to R$.

Definition 3.1. *A g-characteristic is a row vector $m \in \mathbf{Z}^{2g}$. We denote $m = (m'|m'')$ where $m' = (m'_1, m'_2, \cdots, m'_g)$, $m'' = (m''_1, m''_2, \cdots, m''_g) \in \mathbf{Z}^g$. We call the g-chatacteristic m is even (resp. odd) if $\sum_{i=1}^{g} m'_i m''_i$ is even (resp. odd).*

We denote the Siegel upper half space of degree g by $\mathfrak{S}_g$. For a g-characteristic $m = (m'|m'') \in \mathbf{Z}^{2g}$ and $\tau \in \mathfrak{S}_g$, $z \in \mathbf{C}^g$, The theta function θ_m is defined by

$$\theta_m(\tau, z) := \sum_{p \in \mathbf{Z}^g} \exp(\pi i\{(p + m'/2)\tau{}^t(p + m'/2) + (p + m'/2){}^t(z + m''/2)\}).$$

We denote $\theta_m(\tau, 0)$ simply by $\theta_m(\tau)$. Let Ω be the sheaf of holomorphic 1-forms on R. Choose a symplectic basis $\{A_i, B_i\}_{i=1}^{g}$ of $H_1(R; \mathbf{Z})$. It is known that under the homomorphism

$$\begin{array}{ccll} H_1(R; \mathbf{Z}) & \to & H^0(R; \Omega)^* & := \mathrm{Hom}(H^0(R; \Omega), \mathbf{C}), \\ c & \mapsto & (\omega \mapsto \int_c \omega) & \end{array}$$

$H_1(R; \mathbf{Z})$ maps onto a lattice in $H^0(R; \Omega)^*$. The Jacobi variety of R is defined by

$$J(R) = \frac{H^0(R; \Omega)^*}{H_1(R; \mathbf{Z})}.$$

A basis $\{\omega_i\}_{i=1}^{g}$ of $H^0(R; \Omega)$ that satisfies

$$\int_{A_j} \omega_i = \begin{cases} 1, & \text{if } i = j, \\ 0, & \text{if } i \neq j, \end{cases}$$

is called the normalized basis with respect to the symplectic basis $\{A_i, B_i\}_{i=1}^{g}$. For the normalized basis $\{\omega_i\}_{i=1}^{g}$, the g-square matrix

$$\tau = (\tau_{ij}), \qquad \tau_{ij} = \int_{B_j} \omega_i$$

is known to be the elements of the Siegel upper half space $\mathfrak{S}_g$, and is called the period matrix. For an even g-characteristic $m = (m'|m'')$ and the period matrix τ, $\theta_m(\tau)$ is called the Riemann theta constant with m associated with the compact Riemann surface R and $\{A_i, B_i\}_{i=1}^{g}$.

Denote the generator of the deck transformation group of the cover $\hat{R} \to R$ by $t : \hat{R} \to \hat{R}$, the (-1)-eigenspace of $t_* : H_1(\hat{R}; \mathbf{Z}) \to H_1(\hat{R}; \mathbf{Z})$ by

$$H_1(\hat{R}; \mathbf{Z})^- = \{c \in H_1(\hat{R}; \mathbf{Z}) \mid t_*(c) = -c\},$$

and the (-1)-eigenspace of $t^* : H^0(\hat{R}; \Omega) \to H^0(\hat{R}; \Omega)$ by

$$H^0(\hat{R}; \Omega)^- = \{\omega \in H^0(\hat{R}; \Omega) \mid t^*(\omega) = -\omega\}.$$

Under the homomorphism

$$\begin{array}{ccll} H_1(\hat{R}; \mathbf{Z}) & \to & H^0(\hat{R}; \Omega)^* & := \mathrm{Hom}(H^0(\hat{R}; \Omega), \mathbf{C}), \\ c & \mapsto & (\omega \mapsto \int_c \omega) & \end{array}$$

$H_1(\hat{R}; \mathbf{Z})^-$ maps onto a lattice in $(H^0(R; \Omega)^-)^*$.

Definition 3.2. *The Prym variety* $\mathrm{Prym}(\hat{R}, p_2)$ *of the cover* p_2 *is defined by*

$$\mathrm{Prym}(\hat{R}, p_2) = \frac{(H^0(\hat{R}; \Omega)^-)^*}{H_1(\hat{R}; \mathbf{Z})^-} \subset J(\hat{R}).$$

For a symplectic basis $\{A_i, B_i\}_{i=1}^{g}$ of $H_1(R; \mathbf{Z})$, we choose a basis of $H_1(\hat{R}; \mathbf{Z})$ as follows. For $i = 1, 2, \cdots, g-1$, denote the two lifts of A_i by $\hat{A}_i$ and $\hat{A}_{i+g}$, and the two lifts of B_i by $\hat{B}_i$ and $\hat{B}_{i+g}$ such that

$$\hat{A}_i \cdot \hat{B}_i = 1.$$

The lifts of $2A_g$ and B_g are uniquely determined, and denote them by $\hat{A}_g$ and $\hat{B}_g$, respectively. Then, $\{A_i - A_{g+i}, B_i - B_{g+i}\}_{i=1}^{g-1}$ form a basis of $H_1(\hat{R}; \mathbf{Z})^-$. Moreover since the basis $\{\hat{A}_i - \hat{A}_{g+i}, \hat{B}_i - \hat{B}_{g+i}\}_{i=1}^{g-1}$ of $H_1(\hat{R}; \mathbf{Z})^-$ satisfies

$$(\hat{A}_i - \hat{A}_{g+i}) \cdot (\hat{A}_j - \hat{A}_{g+j}) = 0, \ (\hat{B}_i - \hat{B}_{g+i}) \cdot (\hat{B}_j - \hat{B}_{g+j}) = 0$$
$$(\hat{A}_i - \hat{A}_{g+i}) \cdot (\hat{B}_j - \hat{B}_{g+j}) = 2\delta_{i\,j}.$$

Therefore, the action of $\hat{\varphi} \in \hat{\mathcal{M}}_{(g)}(p_2)$ on the basis $\{\hat{A}_i - \hat{A}_{g+i}, \hat{B}_i - \hat{B}_{g+i}\}_{i=1}^{g-1}$ induces the homomorphism

$$\tilde{\iota} : \hat{\mathcal{M}}_{(g)}(p_2) \to \mathrm{Sp}(2g-2; \mathbf{Z}).$$

For the above symplectic basis $\{\hat{A}_i, \hat{B}_i\}_{i=1}^{2g-1}$, choose the normalized basis $\{\hat{\omega}_i\}_{i=1}^{2g-1}$ of $H^0(\hat{R}; \Omega)$, then $\{(\hat{\omega}_i - \hat{\omega}_{g+i})/2\}_{i=1}^{g-1}$ is a basis of $H^0(\hat{R}; \Omega)^-$. It is known that the $(g-1)$-square matrix

$$\tilde{\tau} = (\tilde{\tau}_{ij}), \qquad \tilde{\tau}_{ij} = \int_{\hat{B}_j - \hat{B}_{g+j}} \frac{\hat{\omega}_i - \hat{\omega}_{g+i}}{2}$$

is the element of the Siegel upper half space $\mathfrak{S}_{g-1}$. We call $\tilde{\tau}$ the period matrix of the Prym variety.

Definition 3.3. *For an even* $(g-1)$*-characteristic* $\tilde{m} = (\tilde{m}'|\tilde{m}'')$ *and the period matrix* $\tilde{\tau}$ *of* $\mathrm{Prym}(\hat{R}, p_2)$, $\theta_{\tilde{m}}(\tilde{\tau})$ *is called the Schottky theta constant with* $\tilde{m}$ *associated with the cover* $p_2 : \hat{R} \to R$ *and* $\{\hat{A}_i, \hat{B}_i\}_{i=1}^{2g-1}$.

3.2 Definition of the homomorphism $e : \hat{\mathcal{M}}_g(p_2) \to <\sqrt{-1}>$

In this subsection, we give the definition of the homomorphism $e : \hat{\mathcal{M}}_{(g)}(p_2) \to <\sqrt{-1}>$. Let τ be the period matrix of R, and $\tilde{\tau}$ the period matrix of the cover p_2. Consider the function

$$\Phi^{\tilde{m}}_{m,n}(\tilde{\tau},\tau) = \frac{\tilde{\theta}^2_{\tilde{m}}(\tilde{\tau})}{\theta_m(\tau)\theta_n(\tau)}$$

for even g-characteristics m, n and an even $(g-1)$-characteristic $\tilde{m}$. For a generic Riemann surface and a double unbranched covering space, $\Phi^{\tilde{m}}_{m,n}(\tilde{\tau},\tau)$ is known to be a nonzero complex number (Fay[5]). For a g-square matrix $M = (m_{ij})$, denote the row vector obtained by taking the diagonal entries of M by $M_0 := (m_{11}, m_{22}, \cdots, m_{gg}) \in \mathbf{Z}^g$. For $\sigma = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \in \mathrm{Sp}(2g;\mathbf{Z})$ and a g-characteristic m, we define

$$\sigma \cdot m = m \begin{pmatrix} {}^t\alpha & -{}^t\gamma \\ -{}^t\beta & {}^t\delta \end{pmatrix} + (({}^t\beta\alpha)_0 | ({}^t\delta\gamma)_0) \in \mathbf{Z}^{2g}.$$

Note that this is not an action of $\mathrm{Sp}(2g;\mathbf{Z})$ on $\mathbf{Z}^{2g}$, and that this definition is different from that of Igusa[9]. For $\hat{\varphi} \in \hat{\mathcal{M}}_{(g)}(p_2)$, denote $P_2(\hat{\varphi})$ by $\varphi \in \mathcal{M}_g(p_2)$. For an even $(g-1)$-characteristic $\tilde{m}$, choose the g-characteristics $m = (\tilde{m}', 0|\tilde{m}'', 1)$ and $n = (\tilde{m}', 0|\tilde{m}'', 0)$. Define the map $d_{\tilde{m},(\tilde{\tau},\tau)} : \hat{\mathcal{M}}_{(g)}(p_2) \to \mathbf{C}$ by

$$d_{\tilde{m},(\tilde{\tau},\tau)}(\hat{\varphi}) := \frac{\Phi^{\tilde{m}}_{m,n}(\tilde{\tau},\tau)}{\Phi^{\tilde{\iota}(\hat{\varphi})\cdot\tilde{m}}_{\iota(\varphi)\cdot m,\iota(\varphi)\cdot n}(\tilde{\tau},\tau)}.$$

In the next subsection, we will prove that $d_{\tilde{m}} = d_{\tilde{m},(\tilde{\tau},\tau)}$ is independent of the period matrices $\tilde{\tau}$ and τ, and that the image of $d_{\tilde{m}}$ equals $< -1 >$. For $\sigma = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \in \mathrm{Sp}(2g;\mathbf{Z})$ and $\tau \in \mathfrak{S}_g$, we define the action of $\mathrm{Sp}(2g;\mathbf{Z})$ on $\mathfrak{S}_g$ by

$$\sigma \cdot \tau := (\delta\tau + \gamma)(\beta\tau + \alpha)^{-1}.$$

For $\sigma = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \in \mathrm{Sp}(2g;\mathbf{Z})$, it is known that the theta function has the transformation law (see Igusa[10])

$$\theta_{\sigma\cdot m}(\sigma\cdot\tau) = \gamma_m(\sigma)\det(\beta\tau + \alpha)^{-\frac{1}{2}}\theta_m(\tau),$$

where $\gamma_m(\sigma) \in <\exp(\pi/4)>$ is called the theta multiplier.

Now we can construct a homomorphism $e_{\tilde{m}} : \hat{\mathcal{M}}_{(g)}(p_2) \to <\sqrt{-1}>$ using $d_{\tilde{m}}$ and γ_m. For $\hat{\varphi} \in \hat{\mathcal{M}}_{(g)}(p_2)$ and an even $(g-1)$-characteristic $\tilde{m}$, define the map $e_{\tilde{m}}$ by

$$e_{\tilde{m}}(\hat{\varphi}) := d_{\tilde{m}}(\hat{\varphi}) \frac{\gamma^2_{\tilde{m}}(\tilde{\iota}(\hat{\varphi}))}{\gamma_m(\iota(\varphi))\gamma_n(\iota(\varphi))}.$$

Note that $\gamma^2_m(\iota(\varphi))$ and $\gamma_m(\iota(\varphi))\gamma_n(\iota(\varphi))$ are uniquely determined. We will prove that $e = e_{\tilde{m}}$ is a homomorphism independent of the choice of $\tilde{m}$, and that the image of $e_{\tilde{m}}$ equals $<\sqrt{-1}>$ in the next subsection.

3.3 Proof of the main theorem

In this subsection, we will prove that $e_{\tilde{m}} : \hat{\mathcal{M}}_{(g)}(p_2) \to <\sqrt{-1}>$ is a surjective homomorphism. We also prove that $d_{\tilde{m}} = d_{\tilde{m},(\tilde{\tau},\tau)}$ does not depends on the choice of $(\tilde{\tau}, \tau)$, and that the image of $d_{\tilde{m}}$ equals the cyclic group $<-1>$. For $\hat{\varphi} \in \mathcal{M}_{(g)}(p_2)$, we denote simply $\tilde{\iota}(\hat{\varphi}) \in \mathrm{Sp}(2g-2; \mathbf{Z})$ and $\iota(\varphi) = \iota(P(\hat{\varphi})) \in \Gamma_g(p_2)$ by $\tilde{\sigma}$ and σ, respectively.

To prove that $d_{\tilde{m}}$ only depends on $\hat{\varphi} \in \hat{\mathcal{M}}_{(g)}(p_2)$ and $\tilde{m} \in \mathbf{Z}^{g-1}$, we need the following theorem.

Theorem 3.4 (Farkas, Rauch[4])**.**
For an even $(g-1)$-characteristic $\tilde{m}$, define the g-characteristics $m = (\tilde{m}', 0|\tilde{m}'', 1)$ and $n = (\tilde{m}', 0|\tilde{m}'', 0)$. Then $\Phi^{\tilde{m}}_{m,n}(\tilde{\tau}, \tau)$ does not depend on the choice of $\tilde{m}$.

Define $\pi : \mathbf{Z}^{2g} \to \mathbf{Z}^{2g-2}$ by $\pi(m'|m'') = (m'_1, m'_2, \cdots, m'_{g-1} \mid m''_1, m''_2, \cdots, m''_{g-1})$.

Lemma 3.5. *For an even g-characteristic $\tilde{m}$ and $\hat{\varphi} \in \mathcal{M}_{(g)}(p_2)$,*

$$\tilde{\sigma} \cdot \tilde{m} \equiv \pi(\sigma \cdot m) \mod 2,$$

where $m = (\tilde{m}', 0|\tilde{m}'', 1)$.

Proof. Denote the 1-eigenspace of $H_1(\hat{R}; \mathbf{Q})$ by $H_1(\hat{R}; \mathbf{Q})^+$. Then

$$\{\hat{A}_i + \hat{A}_{g+i},\ \hat{B}_i + \hat{B}_{g+i}\}_{i=1}^{g-1} \cup \{\hat{A}_g,\ 2\hat{B}_g\}$$

is a basis of $H_1(\hat{R}; \mathbf{Q})^+$. The restriction of p_2

$$H_1(\hat{R}; \mathbf{Q})^+ \to H_1(R; \mathbf{Q})$$

maps the basis $\{\hat{A}_i + \hat{A}_{g+i},\ \hat{B}_i + \hat{B}_{g+i}\}_{i=1}^{g-1} \cup \{\hat{A}_g,\ 2\hat{B}_g\} \in H_1(\hat{R}; \mathbf{Q})^+$ to the basis $\{2A_i,\ 2B_i\}_{i=1}^{g} \in H_1(R; \mathbf{Q})$. Since for $i = 1, \cdots, g-1$ we have

$$\varphi_*(2A_i) = \varphi_*(p_2)_*(\hat{A}_i + \hat{A}_{g+i}) = (p_2)_*\hat{\varphi}_*(\hat{A}_i + \hat{A}_{g+i}),\ \varphi(2A_g) = (p_2)_*\hat{\varphi}_*(\hat{A}_g),$$
$$\varphi_*(2B_i) = (p_2)_*\hat{\varphi}_*(\hat{B}_i + \hat{B}_{g+i}),\ \text{and } \varphi_*(2B_g) = (p_2)_*\hat{\varphi}_*(2\hat{B}_g).$$

Hence, the induced homomorphism $\hat{\mathcal{M}}_{(g)}(p_2) \to \mathrm{Sp}(2g; \mathbf{Z})$ by the action of $\hat{\varphi} \in \hat{\mathcal{M}}_{(g)}(p_2)$ on the basis $\{\hat{A}_i + \hat{A}_{g+i},\ \hat{B}_i + \hat{B}_{g+i}\}_{i=1}^{g-1} \cup \{\hat{A}_g,\ 2\hat{B}_g\}$ is equal to $\iota P_2 : \hat{\mathcal{M}}_{(g)}(p_2) \to \Gamma_g(p_2)$. Denote $\tilde{\sigma} \in \mathrm{Sp}(2g-2; \mathbf{Z})$ by

$$\tilde{\sigma} = \begin{pmatrix} \alpha' & \beta' \\ \gamma' & \delta' \end{pmatrix},$$

where $\alpha', \beta', \gamma', \delta' \in M(g-1; \mathbf{Z})$. Since we have

$$\hat{\varphi}_*(\hat{A}_i + \hat{A}_{g+i}) \equiv \hat{\varphi}_*(\hat{A}_i - \hat{A}_{g+i}),\ \text{and } \hat{\varphi}_*(\hat{B}_i + \hat{B}_{g+i}) \equiv \hat{\varphi}_*(\hat{B}_i - \hat{B}_{g+i}) \mod 2,$$

and $\sigma = \iota P_2(\hat{\varphi})$ preserves the homology class $B_g \bmod 2$, σ is written in the form

$$\sigma = \begin{pmatrix} \alpha' & {}^t v_1 & \beta' & 0 \\ 0 & 1 & 0 & 0 \\ \gamma' & {}^t v_2 & \delta' & 0 \\ v_3 & k & v_4 & 1 \end{pmatrix} \bmod 2,$$

where $v_1, v_2, v_3, v_4 \in \mathbf{Z}^{g-1}, k \in \mathbf{Z}$. Then it is easy to see that $\pi(\sigma \cdot m) \equiv \tilde{\sigma} \cdot \tilde{m} \mod 2$. ☐

Lemma 3.6. *For $\hat{\varphi} \in \hat{\mathcal{M}}_{(g)}(p_2)$, the value $d_{\tilde{m}}(\hat{\varphi}) = d_{\tilde{m},(\tilde{\tau},\tau)}(\hat{\varphi})$ does not depend on the choice of $(\tilde{\tau}, \tau)$, and the image of $d_{\tilde{m}}$ equals the cyclic group $< -1 >$. In particular, it does not depend on a complex structure of the cover $p_2 : \hat{R} \to R$.*

Proof. Note that, for any g-characteristic $u = (u'|u''), v = (v'|v'') \in \mathbf{Z}$ we have

$$\theta_{u+2v} = (-1)^{u'v''} \theta_u,$$

by the definition of the theta function. Consider the g-characteristic $v_0 = (0, \cdots, 0, 1|0, \cdots, 0, 0) \in \mathbf{Z}^{2g}$. Since σ preserves the homology class $B_g \bmod 2$, we have

$$\sigma \cdot (m - n) = (m - n) \begin{pmatrix} {}^t\alpha & -{}^t\gamma \\ -{}^t\beta & {}^t\delta \end{pmatrix} \equiv v_0 \begin{pmatrix} {}^t\alpha & -{}^t\gamma \\ -{}^t\beta & {}^t\delta \end{pmatrix} \equiv v_0 \ , and$$

$$(\sigma \cdot m)'_g \equiv (\sigma \cdot n)'_g \equiv (\beta {}^t\alpha)_{gg} \equiv 0 \mod 2.$$

By Lemma 3.5, there exists $v_1, v_2 \in \mathbf{Z}^{2g}$ such that

$$\sigma \cdot m + 2v_1 = ((\tilde{\sigma} \cdot \tilde{m})', 0|(\tilde{\sigma} \cdot \tilde{m})'', k_1), \text{ and } \sigma \cdot n + 2v_2 = ((\tilde{\sigma} \cdot \tilde{m})', 0|(\tilde{\sigma} \cdot \tilde{m})'', k_2),$$

where

$$k_1 = 0 \text{ or } 1, \ k_2 = 0 \text{ or } 1, \text{ and } k_1 + k_2 = 1.$$

Then there exists $p(\tilde{m}, \hat{\varphi}) \in < -1 >$ such that

$$\Phi^{\tilde{\sigma}\cdot\tilde{m}}_{\sigma \cdot m + 2v_1, \sigma \cdot n + 2v_2}(\tilde{\tau}, \tau) = p(\tilde{m}, \hat{\varphi}) \Phi^{\tilde{\sigma}\cdot\tilde{m}}_{\sigma \cdot m, \sigma \cdot n}(\tilde{\tau}, \tau).$$

Note that $p(\tilde{m}, \hat{\varphi})$ does not depend on the choice of $(\tilde{\tau}, \tau)$. By Theorem 3.4, we have

$$\Phi^{\tilde{\sigma}\cdot\tilde{m}}_{\sigma \cdot m + 2v_1, \sigma \cdot n + 2v_2}(\tilde{\tau}, \tau) = \Phi^{\tilde{m}}_{m,n}(\tilde{\tau}, \tau).$$

Hence we have

$$p(\tilde{m}, \hat{\varphi}) = d_{\tilde{m}}(\hat{\varphi}).$$

This proves the lemma. ☐

Consider the action of $\varphi \in \mathcal{M}_g(p_2)$ on the symplectic basis $\{A_i, B_i\}_{i=1}^{g}$. The basis $\{\varphi_* A_i, \varphi_* B_i\}_{i=1}^{g}$ is also a symplectic basis of $H_1(R; \mathbf{Z})$. The corresponding period matrix is

$$\tau' = (\tau'_{ij}), \qquad \tau'_{ij} = \int_{\varphi_* B_j} \omega'_i,$$

where $\{\omega'_i\}_{i=1}^{g}$ is the normalized basis. This is equal to ${}^t\iota(\varphi) \cdot \tau$. Next, Consider the action of $\hat{\varphi} \in \hat{\mathcal{M}}_{(g)}(p_2)$ on the basis $\{\hat{A}_i, \hat{B}_i\}_{i=1}^{2g-1}$ of $H_1(\hat{R}; \mathbf{Z})$. Note that the basis $\{\hat{\varphi}_* \hat{A}_i, \hat{\varphi}_* \hat{B}_i\}_{i=1}^{2g-1}$ is again the lift of $\{\varphi_* A_i, \varphi_* B_i\}_{i=1}^{g}$. The period matrix of $\mathrm{Prym}(\hat{R}, p_2)$ with respect to the basis $\{\hat{\varphi}_*(\hat{A}_i - \hat{A}_{g+i}), \hat{\varphi}_*(\hat{B}_i - \hat{B}_{g+i})\}_{i=1}^{2g-1}$ of $H_1(\hat{R}; \mathbf{Z})$ is

$$\tilde{\tau}' := (\tilde{\tau}'_{ij}), \qquad \tilde{\tau}'_{ij} = \int_{\hat{\varphi}_*(\hat{B}_j - \hat{B}_{g+j})} \frac{\hat{\omega'}_i - \hat{\omega'}_{g+i}}{2},$$

where $\{\hat{\omega'}_i\}_{i=1}^{2g-1}$ is the normalized basis. This is equal to ${}^t\tilde{\iota}(\hat{\varphi}) \cdot \tilde{\tau}$. Hence, ${}^t\iota(\varphi) \cdot \tau$ is also the perod matrix of R, and ${}^t\tilde{\iota}(\hat{\varphi}) \cdot \tilde{\tau}$ is also the period matrix of the cover p_2. This shows that the pair $(\tilde{\sigma} \cdot \tilde{\tau}, \sigma \cdot \tau)$ satisfies the condition of Theorem 3.4 for any $\hat{\varphi} \in \hat{\mathcal{M}}_{(g)}(p_2)$.

Theorem 3.7. *The map $e_{\tilde{m}}$ is a homomorphism, and the image of $e_{\tilde{m}}(\hat{\varphi})$ equals $<\sqrt{-1}>$. Moreover $e(\hat{\varphi}) := e_{\tilde{m}}(\hat{\varphi})$ does not depend on the choice of $\tilde{m}$.*

Proof. For $\hat{\varphi} \in \hat{\mathcal{M}}_{(g)}(p_2)$, denote $\sigma_1 := \sigma = \iota P_2(\hat{\varphi}')$, and $\tilde{\sigma}_1 := \tilde{\sigma} = \tilde{\iota}(\hat{\varphi})$. Similarly, denote $\sigma_2 := \iota P_2(\hat{\varphi}')$, $\tilde{\sigma}_2 := \tilde{\iota}(\hat{\varphi}')$, and $\sigma_3 := \iota P_2(\hat{\varphi}\hat{\varphi}')$, $\tilde{\sigma}_3 := \tilde{\iota}(\hat{\varphi}\hat{\varphi}')$. Write σ_i as

$$\sigma_i = \begin{pmatrix} \alpha_i & \beta_i \\ \gamma_i & \delta_i \end{pmatrix} \qquad \text{for } i = 1, 2, 3.$$

We also denote simply $\tilde{\tau}' := \tilde{\sigma}_2 \cdot \tilde{\tau}$, and $\tau' := \sigma_2 \cdot \tau$. Since the pairs $(\tilde{\sigma}_1\tilde{\sigma}_2 \cdot \tilde{\tau}, \sigma_1\sigma_2 \cdot \tau)$, and $(\tilde{\sigma}_2 \cdot \tilde{\tau}, \sigma_2 \cdot \tau)$ satisfies the condition of Theorem 3.4, we have

$$\begin{aligned}
\frac{1}{d_{\tilde{m}}(\widehat{\varphi\varphi'})} &= \frac{\Phi^{(\tilde{\sigma}_1\tilde{\sigma}_2)\cdot\tilde{m}}_{(\sigma_1\sigma_2)\cdot m, (\sigma_1\sigma_2)\cdot n}(\tilde{\sigma}_1\tilde{\sigma}_2 \cdot \tilde{\tau}, \sigma_1\sigma_2 \cdot \tau)}{\Phi^{\tilde{m}}_{m,n}(\tilde{\sigma}_1\tilde{\sigma}_2 \cdot \tilde{\tau}, \sigma_1\sigma_2 \cdot \tau)} = \frac{\gamma^2_{\tilde{m}}(\tilde{\sigma}_1\tilde{\sigma}_2)}{\gamma_m(\sigma_1\sigma_2)\gamma_n(\sigma_1\sigma_2)} \frac{\det(\tilde{\beta}_3\tilde{\tau} + \tilde{\alpha}_3)^{-1}}{\det(\beta_3\tau + \alpha_3)^{-1}} \frac{\Phi^{\tilde{m}}_{m,n}(\tilde{\tau}, \tau)}{\Phi^{\tilde{m}}_{m,n}(\tilde{\sigma}_1\tilde{\sigma}_2 \cdot \tilde{\tau}, \sigma_1\sigma_2 \cdot \tau)}, \\
\frac{1}{d_{\tilde{m}}(\hat{\varphi})} &= \frac{\Phi^{\tilde{\sigma}_1\cdot\tilde{m}}_{\sigma_1\cdot m, \sigma_1\cdot n}(\tilde{\sigma}_1\tilde{\sigma}_2 \cdot \tilde{\tau}, \sigma_1\sigma_2 \cdot \tau)}{\Phi^{\tilde{m}}_{m,n}(\tilde{\sigma}_1\tilde{\sigma}_2 \cdot \tilde{\tau}, \sigma_1\sigma_2 \cdot \tau)} = \frac{\gamma^2_{\tilde{m}}(\tilde{\sigma}_1)}{\gamma_m(\sigma_1)\gamma_n(\sigma_1)} \frac{\det(\tilde{\beta}_1\tilde{\tau}' + \tilde{\alpha}_1)^{-1}}{\det(\beta_1\tau' + \alpha_1)^{-1}} \frac{\Phi^{\tilde{m}}_{m,n}(\tilde{\sigma}_2 \cdot \tilde{\tau}, \sigma_2 \cdot \tau)}{\Phi^{\tilde{m}}_{m,n}(\tilde{\sigma}_1\tilde{\sigma}_2 \cdot \tilde{\tau}, \sigma_1\sigma_2 \cdot \tau)}, \\
\frac{1}{d_{\tilde{m}}(\hat{\varphi'})} &= \frac{\Phi^{\tilde{\sigma}_2\cdot\tilde{m}}_{\sigma_2\cdot m, \sigma_2\cdot n}(\tilde{\sigma}_2 \cdot \tilde{\tau}, \sigma_2 \cdot \tau)}{\Phi^{\tilde{m}}_{m,n}(\tilde{\sigma}_2 \cdot \tilde{\tau}, \sigma_2 \cdot \tau)} = \frac{\gamma^2_{\tilde{m}}(\tilde{\sigma}_2)}{\gamma_m(\sigma_2)\gamma_n(\sigma_2)} \frac{\det(\tilde{\beta}_2\tilde{\tau} + \tilde{\alpha}_2)^{-1}}{\det(\beta_2\tau + \alpha_2)^{-1}} \frac{\Phi^{\tilde{m}}_{m,n}(\tilde{\tau}, \tau)}{\Phi^{\tilde{m}}_{m,n}(\tilde{\sigma}_2 \cdot \tilde{\tau}, \sigma_2 \cdot \tau)},
\end{aligned}$$

by the definition of $d_{\tilde{m}}(\hat{\varphi})$. It is easy to see that

$$\begin{gathered}
\det(\tilde{\beta}_2\tilde{\tau} + \tilde{\alpha}_2)\det(\tilde{\beta}_1\tilde{\tau}' + \tilde{\alpha}_1) = \det(\tilde{\beta}_3\tilde{\tau} + \tilde{\alpha}_3), \text{ and} \\
\det(\beta_2\tau + \alpha_2)\det(\beta_1\tau' + \alpha_1) = \det(\beta_3\tau + \alpha_3).
\end{gathered}$$

This shows that $e_{\tilde{m}}$ is a homomorphism.

Next, we determine the image of $e_{\tilde{m}}$. There are two lifts in $\hat{\mathcal{M}}_{(g)}(p_2)$ of $a^2 \in \mathcal{M}_g(p_2)$. We denote the lift which fix the homology class $\hat{A}_1$ by $\hat{a} \in \hat{\mathcal{M}}_{(g)}(p_2)$. As we stated in Remark 2.5, $H_1(\hat{\mathcal{M}}_{(g)}(p_2); \mathbf{Z})$ is generated by $\hat{a}$. For $\hat{\varphi} = \hat{a}$, we have $\tilde{\sigma} = \tilde{\iota}(\hat{a}) = I_{2g-2} \in \mathrm{Sp}(2g-2;\mathbf{Z})$, $\sigma = \iota P_2(\hat{a}) = \gamma_{gg} \in \Gamma_g(p_2)$. From Theorem 3 in Igusa[9], for any $\tilde{m} \in \mathbf{Z}^{2(g-1)}$, we have

$$\gamma_m(\sigma)\gamma_n(\sigma) = -\sqrt{-1}, \text{ and } \gamma^2_{\tilde{m}}(\tilde{\sigma}) = 1,$$

so that

$$\frac{\gamma^2_{\tilde{m}}(\tilde{\sigma})}{\gamma_m(\sigma)\gamma_n(\sigma)} = \sqrt{-1}.$$

It is easy to see that $d_{\tilde{m}}(\hat{a}) = 1$. Hence $e_{\tilde{m}}(\hat{a})$ is a generator of the cyclic group $<\sqrt{-1}>$ and is independent of the choice of $\tilde{m}$. □

For $r = 0, 1$, we proved $H_1(\hat{\mathcal{M}}_{(g,r)}(p_2);\mathbf{Z}) \cong \mathbf{Z}/2\mathbf{Z}$ or $\mathbf{Z}/4\mathbf{Z}$ in Section 2. From the above Theorem, we have

$$H_1(\hat{\mathcal{M}}_{(g,r)}(p_2);\mathbf{Z}) \cong \mathbf{Z}/4\mathbf{Z}.$$

Since $\mathcal{M}_{g,1}(p_2)$ is isomorphic to $\hat{\mathcal{M}}_{(g,1)}(p_2)$, we have $H_1(\mathcal{M}_{g,1}(p_2);\mathbf{Z}) \cong \mathbf{Z}/4\mathbf{Z}$. Consider $H_1(\mathcal{M}_g(p_2);\mathbf{Z})$ when genus g is odd. For the deck transformation t, we obtain

$$e(t) = (-1)^{g-1},$$

from Theorem 3 in Igusa[9]. By the Lyndon-Hochschild-Serre spectral sequence, we have

$$\mathbf{Z}/2\mathbf{Z} \to H_1(\hat{\mathcal{M}}_{(g)}(p_2);\mathbf{Z}) \to H_1(\mathcal{M}_g(p_2);\mathbf{Z}) \to 0$$

This shows that $H_1(\mathcal{M}_g(p_2);\mathbf{Z}) \cong \mathbf{Z}/4\mathbf{Z}$ when g is odd. This completes the proof of Theorem 0.2. From the Theorem 0.2, we obtain many homomorphisms $\mathcal{M}_{g,1}[d] \to \mathbf{Z}/4\mathbf{Z}$ for an even integer d.

Proposition 3.8. *For a positive even integer d, there exists an injection*

$$(\mathbf{Z}/4\mathbf{Z})^{2g} \hookrightarrow \mathrm{Hom}(\mathcal{M}_{g,1}[d];\mathbf{Z}/4\mathbf{Z}).$$

When $d = 2$ and g is odd, we have

$$(\mathbf{Z}/4\mathbf{Z})^{2g} \hookrightarrow \mathrm{Hom}(\mathcal{M}_g[d];\mathbf{Z}/4\mathbf{Z}).$$

Proof. To prove the proposition, we will construct a homomorphism from $\mathcal{M}_{g,1}[d]$ into $\mathcal{M}_{dg/2-1,1}(p'_X)$ for a certain double cover p'_X.

Let X be one of the homology classes $A_1, \cdots, A_g, B_1, \cdots, B_g \in H_1(\Sigma_g;\mathbf{Z})$. Consider the d cover $q_X : \Sigma_{dg-1} \to \Sigma_g$ such that the monodromy homomorphisms $\pi_1(\Sigma_g) \to \mathbf{Z}/d\mathbf{Z}$ is equal to the Poincaré dual of $X \in H^1(\Sigma_g;\mathbf{Z}/d\mathbf{Z})$. Denote a generator of the deck transformation group by t_X. Consider

$$\Sigma_{g,1} = \Sigma_g - D^2 \subset \Sigma_g, \text{ and } \Sigma_{dg-1,d} = \Sigma_{dg-1} - q_X^{-1}(D^2).$$

We denote the restriction of the cover $q_X|_{\Sigma_{g,1}} : \Sigma_{dg-1,d} \to \Sigma_{g,1}$ by p_X. Choose two connected components D_1 and D_2 of $q_X^{-1}(D^2)$ such that $t_X^{d/2} D_1 = D_2$. Consider $\Sigma_{dg-1,2} = \Sigma_{dg-1} - \amalg_{i=1}^2 D_i$. Then we have the double cover

$$p'_X : \Sigma_{dg-1,2} \to \Sigma_{dg-1,2}/ < t_X^{d/2} >= \Sigma_{dg/2-1,1}.$$

We have the projection $P_X : \hat{\mathcal{M}}_{(g,1)}(p_X) \to \mathcal{M}_{g,1}(p_X)$ and $P'_X : \hat{\mathcal{M}}_{(dg/2-1,1)}(p'_X) \to \mathcal{M}_{dg/2-1,1}(p'_X)$. Since the centralizer of $< t_X >$ is included in the centralizer of $< t_X^{d/2} >$, we have the homomorphism

$$\begin{array}{rccc} Q_X : & \hat{\mathcal{M}}_{(g,1)}(p_X) & \to & \hat{\mathcal{M}}_{(dg/2-1,1)}(p'_X). \\ & [\hat{f}] & \mapsto & [\hat{f} \cup id_{\cup_{i=1}^{d-2} D^2}] \end{array}$$

Note that we have the inclusion map $i_X : \mathcal{M}_{g,1}[d] \to \mathcal{M}_{g,1}(p_X)$. Hence we have the homomorphism

$$P'_X Q_X P_X^{-1} i_X : \mathcal{M}_{g,1}[d] \to \mathcal{M}_{dg/2-1,1}(p'_X).$$

Consider the induced homomorphism $(P'_X Q_X P_X^{-1} i_X)_* : H_1(\mathcal{M}_{g,1}[d]; \mathbf{Z}) \to H_1((\mathcal{M}_{dg/2-1,1}(p'_X); \mathbf{Z})$. For the simple closed curves $Y = A_1, \cdots, A_g, B_1, \cdots, B_g$, denote the Dehn twists along Y by T_Y. Then we have

$$(P'_X Q_X P_X^{-1} i_X)_*(T_Y^d) = \begin{cases} 1, & \text{if } Y = X, \\ 0, & \text{otherwise,} \end{cases}$$

by Remark 2.5. Hence the induced map

$$(\mathbf{Z}/4\mathbf{Z})^{2g} \to \mathrm{Hom}(\mathcal{M}_{g,1}[d]; \mathbf{Z}/4\mathbf{Z})$$

is injective.

Next, consider the case of $d = 2$ and g is odd. Then $H_1(\mathcal{M}_g(p_X); \mathbf{Z})$ is isomorphic to $\mathbf{Z}/4\mathbf{Z}$. The inclusion $\mathcal{M}_g[2] \to \mathcal{M}_g(p_X)$ induces a homomorphism $H_1(\mathcal{M}_g[2]; \mathbf{Z}) \to H_1(\mathcal{M}_g(p_X); \mathbf{Z}) \cong \mathbf{Z}/4\mathbf{Z}$. Similarly, we have the injective homomorphism $(\mathbf{Z}/4\mathbf{Z})^{2g} \to \mathrm{Hom}(\mathcal{M}_g[2]; \mathbf{Z}/4\mathbf{Z})$. This completes the proof. □

References

[1] J.S. Birman and R. Craggs, *The μ-Invariant of 3-Manifolds and Certain Structural Properties of the Group of Homeomorphisms of a Closed, Oriented 2-Manifold*, Transactions of the American Mathematical Society **237** (1978), 283–309.

[2] J.S. Birman and H.M. Hilden, *On Isotopies of Homeomorphisms of Riemann Surfaces*, The Annals of Mathematics **97** (1973), no. 3, 424–439.

[3] B. Farb, *Some problems on mapping class groups and moduli space*, Arxiv preprint math.GT/0606432 (2006).

[4] H.M. Farkas and H.E. Rauch, *Period Relations of Schottky Type on Riemann Surfaces*, The Annals of Mathematics **92** (1970), no. 3, 434–461.

[5] J.D. Fay, *Theta functions on Riemann surfaces*, Springer, 1973.

[6] R. Hain, *Torelli groups and Geometry of Moduli Spaces of Curves*, Current Topics in Complex Algebraic Geometry (CH Clemens and J. Kollar, eds.) MSRI publications **28** (1995), 97–143.

[7] J. L. Harer, *The second homology group of the mapping class group of an orientable surface*, Inventiones Mathematicae **72** (1983), no. 2, 221–239.

[8] ______, *The rational Picard group of the moduli space of Riemann surfaces with spin structure*, Contemp. Math **150** (1993), 107–136.

[9] J. Igusa, *On the Graded Ring of Theta-Constants*, American Journal of Mathematics **86** (1964), no. 1, 219–246.

[10] ______, *Theta functions*, Springer, 1972.

[11] D. Johnson, *Homeomorphisms of a Surface which Act Trivially on Homology*, Proceedings of the American Mathematical Society **75** (1979), no. 1, 119–125.

[12] ______, *An abelian quotient of the mapping class group* $\mathcal{I}_g$, Mathematische Annalen **249** (1980), no. 3, 225–242.

[13] ______, *Quadratic Forms and the Birman-Craggs Homomorphisms*, Transactions of the American Mathematical Society **261** (1980), no. 1, 235–254.

[14] ______, *The structure of the Torelli Group III: The abelianization of* $\mathcal{I}_g$, Topology **24** (1985), no. 2, 127–144.

[15] R. Lee, E. Miller, and S. Weintraub, *The Rochlin invariant, theta functions and the holonomy of some determinant line bundle*, J. reine angew. Math **392** (1988), 187–218.

[16] J.D. McCarthy, *On the first cohomology group of cofinite subgroups in surface mapping class groups*, Topology **40** (2000), no. 2, 401–418.

[17] J. Powell, *Two Theorems on the Mapping Class Group of a Surface*, Proceedings of the American Mathematical Society **68** (1978), no. 3, 347–350.